

Post Natal Depression

About this leaflet

This leaflet is for anyone who suffers from postnatal depression (PND for short). We hope it will also be helpful to family and friends and to anyone who wants to know more about this problem. The leaflet describes what it's like to have PND, ways of helping yourself and some of the treatments available. There are references to research information and suggestions for further reading and help.

What is PND?

Postnatal Depression is what happens when you become depressed after having a baby. Sometimes, there may be an obvious reason, often there is none. It can be particularly distressing when you have so looked forward to having your baby through the months of pregnancy. You may feel guilty for feeling like this, or even feel that you can't cope with being a mother.

How common is it?

Around one in every ten women have PND after having a baby. If untreated, it can last for months, or sometimes longer.

What does it feel like to have PND?

Depressed

You feel low, unhappy and wretched for much or all of the time. You may feel worse at particular times of the day, like mornings or evenings. Sometimes, there are good days that make you hope that it is over. It can be very disappointing when they are followed by bad days. Sometimes, it can seem that life is not worth living.



Irritable

You may get irritable with other children and, occasionally, with your baby. You are most likely to get 'ratty' with your partner, who may well wonder what is wrong.

Tired

All new mothers get pretty weary, but depression can make you feel so utterly exhausted that you feel physically ill.

Sleepless

When, at last, you get to bed you find you can't fall asleep. You wake at the crack of dawn, even if your partner has fed the baby overnight.

Not hungry

Depressed mothers usually haven't the time or the interest to eat, and this can make you feel irritable and run down. On the other hand, if you find yourself eating for comfort, you may feel guilty and uncomfortable about getting fat.

Unable to enjoy anything

You find that you can't enjoy or be interested in anything. This may be especially true of sex. Some women get interested in sex again before the 6 week postnatal check-up, but PND usually takes away any desire or enthusiasm. Your partner may seek the comfort and excitement of intercourse, but you don't. This can put a further strain on the relationship.

There are, of course, many other reasons for you to lose interest in sex after having a baby - it may be painful, you may be too tired, or you may be just trying to adjust to having a child.

Unable to cope

PND can make you feel that you have too little time, do nothing well, and that you can't do anything about it. It can be hard to establish a new routine to cope with the baby, as well as everything else.

Anxious

You may find that you are afraid to be alone with your baby. You may worry that he or she might scream, or choke, or be harmed in some way. Instead of feeling close to your baby, you may feel detached. You can't work out what your baby is feeling, or what your baby needs.

You may feel anxious even if you have strong loving feelings for your baby. You worry that you might lose him or her through infection, mishandling, faulty development or a 'cot death'. You worry about 'snuffles', or how much weight has been (or not been) gained. You worry if your baby is crying or is too quiet (has the baby stopped breathing?) You may find that you need reassurance all the time from your partner, the health visitor, the GP, your family or a neighbour.

You may also worry about your own health. You may feel panicky - your pulse races, your heart thumps and you may feel that you have heart disease or are on the brink of a stroke. Your tiredness may make you wonder if you have some dreadful illness, or if you will ever have any energy again.

The fear of being left alone with all this can cause even the most capable person to cling desperately to their partner, not wanting to be left alone.

Doesn't everybody get depressed after having a baby?

No. About half the women who give birth feel a bit weepy, flat and unsure of themselves on the third or fourth day after having a baby. This is known as the 'Baby Blues', and it passes after a few days.

When does PND happen?

Most cases of PND start within a month of the birth, but it can start up to six months later.

What causes PND?

We don't know enough about why women get PND to be sure who will or won't suffer from it. There is probably no single reason, but a number of different stresses may add up to cause it. We know you are more likely to have PND if you:

- have had depression (especially PND) before
- do not have a supportive partner
- have a premature or sick baby
- lost your own mother when you were a child
- have experienced several stresses in a short period of time. These could be things like a bereavement, you or your partner losing a job, or housing and money problems.

In spite of this, you can still suffer from PND when none of these things have happened and there is no obvious reason.

What about hormones?

In some cases, PND may have something to do with the huge hormone changes which take place at the time of giving birth. Levels of oestrogen, progesterone (and other hormones to do with conception and birth) drop suddenly after the baby is born. How exactly they affect your mood and emotions is not clear. No real differences have been found in the hormone changes of women who do and do not get PND. It may be that some women are more sensitive to these changes than others.

Do women with PND harm their babies?

Very rarely. It is more likely (although still uncommon) if a mother has a more severe mental disorder, 'puerperal psychosis'. This is a very serious (but treatable) mental illness which usually comes on within a few days of the birth. The mother may become deluded and feel that her baby is evil. Rarely, she may feel so suicidal that she decides to take the baby's life with her own. Puerperal psychosis is much less common than PND: it occurs only after one birth in 500. It is fortunately very rare for a mother to kill her baby.

Occasionally, through utter tiredness and desperation, you might feel like hitting or shaking your baby. Many mothers (and fathers) occasionally feel like this, not just those with PND. In spite of having these feelings at times, most mothers never act on them. This is the case in PND. The problem is more likely to be you worry that you might harm your baby, rather than actually doing so.

What can be done?

A great deal, but first the depression must be recognised. In the past it has often been overlooked or dismissed as the 'baby blues'.

Many depressed mothers don't realise what is wrong with them. They feel ashamed to admit that they are less than thrilled by being a mother. They may think that if they say how they feel, then their baby may be taken away (this won't happen).



Now that there is a greater awareness of depression in general, PND should be missed less often. A questionnaire, such as the Edinburgh Postnatal Depression Scale, can sometimes be used. This can help health visitors and GPs to spot PND.

Ways of helping yourself

Say how you feel:


If you feel miserable, irritable, incompetent, frightened and not all that keen on your baby, then tell someone. Many other women have gone through the same experience. If you don't feel you can talk to your family or friends, talk to your health visitor or GP. They will know that these feelings are common and will be able to help.

Don't be frightened by the diagnosis: It can actually be a relief when someone tells you that you have PND. At last you know what is wrong, and that many others have had the same problem. You can be reassured that you will get better in time. It can help your partner, friends and family to know this as well. They will feel more able to help when they know what the problem is.

Ways for other people to help

Don't be shocked or disappointed if your wife, partner, sister or girlfriend reveals that she has felt awful since the birth of her baby. Take time to listen sympathetically and make sure that she gets the help she needs.

Try not to be shocked or disappointed by a diagnosis of PND. In a way, it is good news because we know it can be effectively helped.



"JANE DARLING,
ISN'T THAT THE BABY CRYING FOR YOU AGAIN?"

Do all you can to help with the practical things that need to be done, while your partner does not feel up to doing them - shopping, feeding and changing the baby, or housework. It may be difficult for a while, but it is worth it.

Make sure that you are clear about what is happening and that you get advice on how to help, especially if you are the mother's partner. Make sure that you have some support yourself. If this is your first baby, you may feel pushed to one side, both by the baby and by your partner's needs. Try not to feel resentful. Your partner needs your support and encouragement. Practical help with the baby, sympathetic listening, patience, affection and being positive will go a long way. Your partner will appreciate this even when the depression is over.

What if I don't want treatment?

Most women will get better without any treatment after a period of weeks, months or sometimes longer. However, this can mean a lot of suffering. PND spoils the experience of new motherhood, and will strain your relationship with your baby and partner. So the shorter it lasts, the better. It is advisable to diagnose and treat PND as soon as possible.

What about talking treatments?

It can be a great relief just to talk to a sympathetic, understanding, uncritical listener - this could be a friend, a relative, a volunteer or a professional. Many general practices now have a counsellor, and trained health visitors can help treat PND.

There are more specialised psychological treatments. **Psychotherapy** can help you to understand the depression in terms of what has happened to you in the past. **Cognitive therapy** can help you to understand the depression in terms of how you think about yourself, the world and other people. These can be arranged through your GP with a community psychiatric nurse, a psychologist or a psychiatrist.

Are there problems with these treatments?

These treatments are usually very safe, but they can have unwanted effects. Talking about things may bring up bad memories from the past and this can make you low or distressed. Other people have found that therapy puts a strain on their relationship with their partner.

It is important to make sure that you can trust your therapist and that they have the necessary training. Another problem with talking therapies is that they are still scarce in some areas. There are long waiting lists, so you may not get any treatment for quite a while.

What about tablets?

If you have a more severe depression, or it has not improved with support and reassurance, one of the antidepressant drugs will probably help. Antidepressants take two weeks or so to start working and should be taken for four to six months after you start to feel better.

How do they work?

Antidepressants affect the activity of two chemicals in the brain, Serotonin (also called 5HT) and Noradrenaline.

Do antidepressants have side effects?

Modern antidepressants are safe. They may cause nausea or an increase in anxiety in the early stages, but these usually wear off. Others can make you sleepy or give you a dry mouth. Make sure that your doctor knows that you are breast-feeding. It should be safe to continue breast-feeding as there are antidepressants which do not pass into breast milk to any significant degree, so your baby should not be affected by them. Some people get mild withdrawal symptoms when they stop these medicines, so it's best to come off them slowly. For more information, see our factsheet on antidepressants.

Hormones have been suggested as a treatment for PND. However, there is little evidence that they work, and they have their own dangers, particularly if you have had thromboses (blood clots in the veins) of any sort.

So which treatment is best?

Everyone can try the simple measures outlined in this leaflet. Talking treatments and antidepressants are equally effective, but antidepressants are more likely to be recommended if the depression is severe or has gone on for a long time. They also work a bit quicker than talking treatments. Talking treatments and antidepressants can be given together. Your GP or health visitor will be willing to give advice. It is also sometimes helpful to talk over the options with your family or a close friend. It is important that you feel comfortable with the choice of help or treatment.

Mothers with special needs

Mothers who have a previous history of mental health problems or physical or learning disability do face additional problems, or, equally challenging, additional scrutiny. They should get whatever help and reassurance they need to maintain their own well-being and that of their baby.

Self-help

We don't yet know enough about PND to prevent it in the first place, but certain principles make sense:

DON'T try to be 'superwoman'. Try to do less during your pregnancy and make sure that you don't over-tire yourself. If you are at work, make sure you get regular meals and put your feet up in the lunch hour.

DON'T move house (if you can help it) while you are pregnant or until the baby is six months old.

DO make friends with other couples who are expecting or have just had a baby; among other things, this could lead to a baby-sitting circle.

DO find someone you can talk to. It helps so much to have a close friend you can turn to. (If you can't easily find someone, try the National Childbirth Trust or MAMA - their local groups are very supportive both before and after childbirth).

DO go to ante-natal classes - and take your partner with you.

DO keep in touch with your GP and your health visitor if you have suffered PND before. Any signs of PND can be recognised early and you can start treatment at once. After the baby has arrived:

DO take every opportunity to get your head down. Try to learn to cat-nap. Your partner can give the baby a bottle-feed at night. If you like, you can use your own expressed breast milk for this.

DO get enough nourishment. Healthy foods like salads, fresh vegetables, fruit, fruit juices, milk and cereals are all nice, packed with vitamins and don't need much cooking.

DO find time to have fun with your partner. Try to find a baby-sitter and get out together for a meal, a show or to see friends.

DO let yourself and your partner be intimate if you can: at least kiss and cuddle, stroke and fondle. This will comfort you both and lead to the return of full sexual feelings sooner. Do not feel guilty if this takes some time.

DON'T blame yourself or your partner: life is tough at this time, and tiredness and irritability on both sides can lead to quarrels. 'Having a go' at each other may weaken your relationship when it needs to be at its strongest.

DON'T be afraid to ask for help when you need it. If you have learnt about PND from ante-natal classes (and leaflets like this), you may spot the warning signs before anyone else.

FINALLY, even if you have been depressed for a while, support, counselling and medication can all help you to get better. It's never too late.

Other useful leaflets produced by the Royal College of Psychiatrists are: Alcohol and Depression, Depression, Depression in Older Adults, Manic Depression, Men Behaving Sadly and Physical Illness and Mental Health. We also have factsheets on Antidepressants, and Cognitive Therapy. These can all be downloaded from our website: www.rcpsych.ac.uk. For a catalogue of our materials for the general public, contact the Leaflets Department, Royal College of Psychiatrists, 17 Belgrave Square, London SW1X 8PG. Tel: 020 7235 2351 ext.259; Fax: 020 7235 1935; E-mail: leaflets@rcpsych.ac.uk.

References

Antidepressant treatment for post-natal depression (Cochrane Review) Hoffbrand S, Howard L, Crawley H (2001). Compares antidepressant treatment and counseling. (www.update-software.com)

Depression in postpartum and non-postpartum women: prevalence and risk factors. Eberhard G A et al. (2002) Acta Psychiatrica Scandinavica, Vol 106, 426-433

Controlled trial of the short- and long-term effect of psychological treatment of postpartum depression 2. Impact on the mother-child relationship and child outcome Murray L, Cooper PJ et al (2003) The British Journal of Psychiatry, Vol 182: 420-427

Edinburgh Postnatal Depression Scale (EPDS) British Journal of Psychiatry (1987), Vol. 150 by Cox J L, Holden J M, Sagovsky R (www.wellmother.com/articles/edinburgh.htm)

Can we identify mothers at risk for postpartum depression in the immediate postpartum period using the Edinburgh Postnatal Depression Scale? Dennis C L (2004) Journal of Affective Disorders, Vol. 78, no. 2, p. 163-169

Oestrogens and progestogens for preventing and treating postnatal depression (Cochrane Review). Lawrie T A, Herxheimer A, Dalton K in: The Cochrane Library, Issue 1, 2004. Chichester, UK: John Wiley & Sons, Ltd.

Organisations that can help

Association for Postnatal Depression 145 Dawes Road, Fulham, London SW6 7EB.
Helpline: 020 7386 0868; www.apni.org. Provides support to mothers suffering from post-natal illness. It exists to increase public awareness of the illness and to encourage research into its cause and nature.

CRY-SIS, BM-CRY-SIS, London WC1N 3XX. Tel: 020 7404 5011 (line open 9.00 am to 10.00 pm, 365 days a year); www.cry-sis.com/index.asp Provides self-help and support for families with excessively crying and sleepless babies.

Meet-A-Mum-Association (MAMA), 376 Bideford Green, Linslade, Leighton Buzzard, Beds LU7 2TY. Tel: 01525 21704; E-mail: Meet-A-Mum.Assoc@blueyonder.co.uk;
www.mama.org.uk; Helpline: 020 8768 0123 (7.00 pm to 10.00 pm weekdays) Self-help groups for mothers with small children and specific help and support to women suffering from postnatal depression.

National Childbirth Trust, Alexandra House, Oldham Terrace, Acton, London W3 6NH.
Enquiry line: 0870 444 8707; Breastfeeding line: 0870 444 8708; E-fax: 0870 770 3237;
www.nctpregnancyandbabycare.com. Advice, support and counselling on all aspects of childbirth and early parenthood.

The Samaritans. Tel: 08457 909090 (UK) or 1850 609090 (Eire); E-mail: jo@samaritans.org;
www.samaritans.org. Provides confidential emotional support to any person who is suicidal or despairing.

Books

Feelings after birth: The NCT Book of Postnatal Depression by Healthier Welford (Book Production Consultants)

Surviving Post-natal Depression: At home no-one hears you scream by Cara Aitken (Jessica Kingsley Publishers)

Depression after childbirth: How to recognise, treat and prevent postnatal depression by Katherina Dalton (Oxford University Press)

Copyright: Copyright: © Royal College of Psychiatrists, 2004, all rights reserved.

Registered charity number 228636

Last updated May 2004 Produced by the Royal College of Psychiatrists' Public Education Committee Editorial Sub-Committee Series Editor: Dr Philip Timms User & Carer input: Royal College of Psychiatrists' Special Committee of Patients and Carers Expert Review: Dr R. Cantwell Illustrations by Professor Tom Barnes

This leaflet may not be reproduced in whole or in part without the permission of the Royal College of Psychiatrists. Royal College of Psychiatrists. Registered charity number 228636